

Research Article

Relatively Bounded Perturbations of a Closed Linear Relation and Its Adjoint

Guixin Xu 

School of Mathematics and Statistics, Beijing Technology and Business University, Beijing, China

Correspondence should be addressed to Guixin Xu; guixinxu_ds@163.com

Received 10 February 2022; Revised 14 April 2022; Accepted 3 May 2022; Published 25 May 2022

Academic Editor: Marek Galewski

Copyright © 2022 Guixin Xu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with the relatively bounded perturbations of a closed linear relation and its adjoint in Hilbert spaces. A stability result about orthogonal projections onto the ranges of linear relations is obtained. By using this result, two perturbation theorems for a closed relation and its adjoint are given. These results generalize the corresponding results for single-valued linear operators to linear relations and some of which weaken certain assumptions of the related existing results.

1. Introduction

Perturbation theory is one of the main topics in both pure and applied mathematics. In particular, the perturbations of linear operators (i.e., single-valued operators) have received lots of attention and many useful results have been obtained (cf. [1–4]).

However, when considering the adjoint of a nondensely defined linear operator and the minimal and maximal operators corresponding to a linear discrete Hamiltonian system or a linear symmetric difference equation (cf. [5, 6]), the classical perturbation theory of linear operator is not available in these cases. So, we should apply the perturbation theory of multivalued linear operators to study the above problems. Further, multivalued linear operator theory may provide some useful tools for the study of some Cauchy problems associated with parabolic type equations in Banach spaces [7] and boundary value problems for differential operators [8]. Due to these reasons, it is necessary and urgent to study some topics about multivalued linear operators, which are a necessary foundation of research on those related problems about differential or difference operators.

Note that the graph $G(T)$ of a linear operator or multivalued linear operator T from a linear space X to a linear space Y is a linear subspace in the product space $X \times Y$. Further, it is more convenient to introduce concepts of the

inverse, closure, and adjoint for linear subspaces. So, we shall directly study linear subspace (briefly, subspace) in the product space $X \times Y$. A subspace is also called a linear relation (briefly, relation). A linear operator always means a single-valued linear operator for convenience in the present paper.

To the best of our knowledge, linear relations were introduced by von Neumann [9], motivated by the need to consider adjoint operators of nondensely defined linear differential operators. The operational calculus of linear relations was developed by Arens [10]. His works were followed by many scholars, and some basic concepts, fundamental properties, self-adjoint extension, resolvent, spectrum, and perturbation for linear relations were studied (cf. [5–8, 11–26]).

There are still many important fundamental problems about linear relations that have neither been studied nor completed. It is shown that the closedness and self-adjointness of linear relation are stable under relatively bounded perturbation (cf. [22, 25]). However, they have not been specifically and thoroughly studied. In the present paper, enlightened by the methods used in [4], we shall deeply study the stability of a closed linear relation and its adjoint under more general relatively bounded perturbation in Hilbert spaces. The spaces X and Y are always assumed to be Hilbert spaces throughout the present paper. The results

obtained in the present paper not only cover the related existing results about operators but also some of them weaken the conditions of the corresponding existing results (see Remarks 1 and 2).

The rest of this paper is organized as follows. In Section 2, some notations, basic concepts, and fundamental results about linear relations are introduced. In Section 3, we first give in Theorem 1 a stability result about orthogonal projections onto the ranges of linear relations, which generalize the corresponding result ([4], Theorem 5.25) for linear operators to linear relations. Then, we investigate the relatively bounded perturbations of a closed linear relation and its adjoint. It is shown that the adjoint of the sum of two linear relations is equal to the sum of each adjoint (see Theorem 3).

2. Preliminaries

In this section, we shall introduce some basic concepts and give some fundamental results about linear relations, which will be used in the sequent sections.

Let X and Y be Hilbert spaces over the complex field C . The norm of $X \times Y$ is defined by

$$\|(x, y)\| = (\|x\|_X^2 + \|y\|_Y^2)^{1/2}, \quad x \in X, y \in Y, \quad (1)$$

where $\|\cdot\|_X$ and $\|\cdot\|_Y$ are the norms of the spaces X and Y , respectively, still denoted by $\|\cdot\|$ without any confusion. The inner product of $X \times Y$ is defined by

$$\begin{aligned} \langle (x_1, y_1), (x_2, y_2) \rangle &= \langle x_1, x_2 \rangle + \langle y_1, y_2 \rangle, \\ (x_1, y_1), (x_2, y_2) &\in X \times Y. \end{aligned} \quad (2)$$

Let $E \subset X$, E^\perp denotes the orthogonal complement of E .

Any linear subspace $T \subset X \times Y$ is called a *linear relation* (briefly, *relation* or *subspace*) of $X \times Y$. $LR(X, Y)$ denotes the set of all linear relations of $X \times Y$. In the case that $X = Y$, $LR(X)$ denotes $LR(X, X)$ briefly.

The domain $D(T)$, range $R(T)$, and null space $N(T)$ of T are, respectively, defined by

$$\begin{aligned} D(T) &:= \{x \in X: (x, y) \in T \text{ for some } y \in Y\}, \\ R(T) &:= \{y \in Y: (x, y) \in T \text{ for some } x \in X\}, \\ N(T) &:= \{x \in X: (x, 0) \in T\}. \end{aligned} \quad (3)$$

A linear relation T is said to be injective if $N(T) = 0$ and surjective if $R(T) = Y$. Further, denote

$$\begin{aligned} T(x) &:= \{y \in Y: (x, y) \in T\}, \\ T^{-1} &:= \{(y, x): (x, y) \in T\}. \end{aligned} \quad (4)$$

It is evident that $T(0) = \{0\}$ if and only if T can uniquely determine a linear operator from $D(T)$ into Y whose graph is T . For convenience, a linear operator (i.e., single-valued operator) from X to Y will always be identified with a relation in $X \times Y$ via its graph. In addition, $N(T) = \{0\}$ if and only if $T^{-1}(0) = \{0\}$, i.e., T is injective if and only if T^{-1} is a linear operator. Further, T is said to be *closed* if $T = \overline{T}$, where \overline{T} is the closure of T .

Let $T, A \in LR(X, Y)$ and $\alpha \in C$. Define

$$\begin{aligned} \alpha T &:= \{(x, \alpha y): (x, y) \in T\}, \\ T + A &:= \{(x, y + z): (x, y) \in T, (x, z) \in A\}, \\ T - \alpha &:= \{(x, y - \alpha x): (x, y) \in T\} \text{ in the case that } Y = X. \end{aligned} \quad (5)$$

If $T \cap A = \{(0, 0)\}$, denote

$$T \dot{+} A := \{(x_1 + x_2, y_1 + y_2): (x_1, y_1) \in T, (x_2, y_2) \in A\}. \quad (6)$$

Further, if T and A are orthogonal, that is, $\langle (x, y), (u, v) \rangle = 0$ for all $(x, y) \in T$ and $(u, v) \in A$, then denote

$$T \oplus A := T \dot{+} A. \quad (7)$$

The *product* of linear relations $T \in LR(X, Y)$ and $A \in LR(Y, Z)$ is defined as follows (see [10]):

$$\begin{aligned} AT &:= \{(x, z) \in X \times Z: (x, y) \in T, \\ &\quad (y, z) \in A \text{ for some } y \in Y\}. \end{aligned} \quad (8)$$

Note that if A and T are operators, then AT is also an operator.

Definition 1. Let $T \in LR(X, Y)$. The *adjoint* T^* of T is defined as a relation from Y to X by

$$T^* := \{(f, g) \in Y \times X: \langle g, x \rangle = \langle f, y \rangle \text{ for all } (x, y) \in T\}. \quad (9)$$

T is said to be *Hermitian* in X^2 if $T \subset T^*$ and said to be *self-adjoint* in X^2 if $T = T^*$

Let U be the flip-flop operator from $X \times Y$ to $Y \times X$ defined by

$$U(x, y) = (y, -x), \quad (x, y) \in X \times Y. \quad (10)$$

It is clear from Definition 1 that

$$T^* = (UT) = UT, \quad (11)$$

where the orthogonal complements refer to the component wise inner product in $Y \times X$ and $X \times Y$, respectively.

Next, we shall briefly recall the concepts of bounded and relatively bounded relations, which were introduced in [17, 22].

Let $T \in LR(X, Y)$. By Q_T , or simply Q , when there is no ambiguity about the relation T , denote the natural quotient map from Y onto $Y/\overline{T(0)}$. Clearly, QT is an operator [17]. Further, denote $B_X := \{x \in X: \|x\| \leq 1\}$.

Definition 2. Let $T \in LR(X, Y)$. For any given $x \in D(T)$, the *norms* of $T(x)$ and T are defined by

$$\begin{aligned} \|T(x)\| &:= \|(QT)(x)\|, \|T\| := \|QT\| \\ &= \sup\{\|(QT)(x)\|: x \in D(T) \cap B_X\}, \end{aligned} \quad (12)$$

respectively. If $\|T\| < \infty$, T is said to be *bounded*.

Definition 3. Let T and A be two linear relations in $X \times Y$ with $D(T) \subset D(A)$. The linear relation A is said to be T -*bounded* if there exist nonnegative numbers a and b such that

$$\|A(x)\| \leq a\|x\| + b\|T(x)\|, \quad x \in D(T). \quad (13)$$

If A is T -*bounded*, then the infimum of all numbers $b \geq 0$ for which (13) holds with some constant $a \geq 0$ is called the T -*bound* of A .

Lemma 1 (Propositions II.1.4 and II.1.5 in [17]). *Let T and S be two linear relations in $X \times Y$. Then,*

- (i) $\|T(x)\| = d(y, T(0)) = d(0, T(x))$ for every $x \in D(T)$ and $y \in T(x)$
- (ii) $\|T(x) + S(x)\| \leq \|T(x)\| + \|S(x)\|$ for every $x \in D(T + S)$
- (iii) $\|\alpha T(x)\| = |\alpha| \|T(x)\|$ for $\alpha \in \mathbb{C}$ and $x \in D(T)$

Note that the norm $\|T(x)\|$ is not a real norm since the following inequality may not hold in general (see Exercise II.1.12 in [17]):

$$\|(S - T)(x)\| \geq \|S(x)\| - \|T(x)\|, \quad x \in D(S) \cap D(T). \quad (14)$$

However, it holds under some conditions.

Lemma 2 (Theorem 2.3 in [22]). *Let $T, S \in LR(X, Y)$ satisfy that $D(S) \subset D(T)$ and $\overline{T(0)} \subset \overline{S(0)}$. Then,*

$$\|(S - T)(x)\| \geq \|S(x)\| - \|T(x)\|, \quad x \in D(S). \quad (15)$$

3. Main Results

In this section, we shall investigate the relatively bounded perturbations of a closed linear relation and its adjoint. For this purpose, we need to discuss the stability of orthogonal projections onto the ranges of linear relations.

We first give the following auxiliary results about linear operators.

Lemma 3 (Theorem 4.3 in [4]). *Let T be an operator from X to Y and M be a subspace of Y satisfying that $R(T) \subset \overline{M}$. Then,*

$$\|T(x)\| = \sup\{|\langle Tx, y \rangle| : y \in M \text{ with } \|y\| = 1\}, \quad x \in D(T). \quad (16)$$

Lemma 4 (Theorem 4.33 in [4]). *Let P_1 and P_2 be two orthogonal projections acting on X . Then, we have*

$$\|P_1 - P_2\| = \max\{\rho_{12}, \rho_{21}\}, \quad (17)$$

where

$$\rho_{j,k} = \sup\{\|P_j h\| : h \in R(P_k), \|h\| \leq 1\}. \quad (18)$$

Theorem 1. *Let $A, B \in LR(X, Y)$ satisfying $D(A) \subset D(B)$ and $B(0) \subset A(0)$. Assume that there exists a constant $c \geq 0$ such that*

$$\|B(x)\| \leq c\|A(x)\| \text{ for } x \in D(A). \quad (19)$$

For every $k \in \mathbb{C}$, let P_k denote the orthogonal projection onto $\overline{R(A + kB)}$. Then, $\|P_k - P_0\| \rightarrow 0$ as $k \rightarrow 0$.

Proof. Suppose that $D(A) \subset D(B)$ and $B(0) \subset A(0)$. It follows from (Proposition 2.1 in [22]) that

$$A = A + kB - kB, \quad (20)$$

for every $k \in \mathbb{C}$. □

Case 1. ($c > 0$). Let $0 < |k| < (1/2c)$. For any $x \in D(A)$, it follows from (19), (20), and Lemma 1 that

$$\begin{aligned} \|B(x)\| &\leq c\|(A + kB - kB)(x)\| \\ &\leq c\|(A + kB)(x)\| + c|k|\|B(x)\| \end{aligned} \quad (21)$$

$$\leq c\|(A + kB)(x)\| + \left(\frac{1}{2}\right)\|B(x)\|.$$

Thus,

$$\|B(x)\| \leq 2c\|(A + kB)(x)\|. \quad (22)$$

Let $Y = \overline{R(A + kB) \oplus R(A + kB)}$. Given any $h \in R(P_0) = R(A)$. It can be decomposed as $h = h_1 + h_2$, where $h_1 \in \overline{R(A + kB)}$ and $h_2 \in \overline{R(A + kB)}$. Let $(x, g) \in A + kB$ with $\|g\| = 1$. There exist $g_1, g_2 \in Y$ such that $(x, g_1) \in A$, $(x, g_2) \in B$, and $g = g_1 + kg_2$. By (i) of Lemma 1, we have that for any $\epsilon > 0$, there is $w \in B(x)$ such that

$$\|w\| < \|B(x)\| + \epsilon. \quad (23)$$

This together with Lemma 1, (22), and the fact that $g \in R(A + kB)$ implies that

$$\begin{aligned} \|w\| &< 2c\|(A + kB)(x)\| + \epsilon \\ &\leq 2c\|g\| + \epsilon = 2c + \epsilon. \end{aligned} \quad (24)$$

Since $(x, w), (x, g_2) \in B$. We have $(0, g_2 - w) \in B$. Hence, $(0, g_2 - w) \in A$ by the assumption that $B(0) \subset A(0)$. Consequently, $(0, k(g_2 - w)) \in A$. Further, noting that $(x, g_1) \in A$, we can get that $(x, g_1 + k(g_2 - w)) \in A$. Then,

$$\begin{aligned} \langle P_k h, g \rangle &= \langle h_1, g \rangle = \langle h, g \rangle = \langle h, g_1 + kg_2 \rangle \\ &= \langle h, g_1 + k(g_2 - w) + kw \rangle = \langle h, kw \rangle = \overline{k} \langle h, w \rangle. \end{aligned} \quad (25)$$

It follows from (24) that

$$|\langle P_k h, g \rangle| \leq |k| \|h\| \|w\| < (2c + \epsilon) |k| \|h\|. \quad (26)$$

Therefore,

$$\|P_k h\| \leq 2c|k| \|h\|, \quad (27)$$

by the arbitrariness of ϵ and Lemma 3.

On the other hand, for any $h' \in R(P_k) = R(A + kB)$, we can prove in a completely analogous way that

$$\|P_0 h'\| \leq c|k| \|h'\|, \quad (28)$$

which together with (27) and Lemma 4 yields that $\|P_k - P_0\| \leq 2c|k|$. Therefore, $\|P_k - P_0\| \rightarrow 0$ as $k \rightarrow 0$.

Case 2. ($c = 0$). In this case, condition (19) turns into $\|B(x)\| = 0$. Then, $\|w\| < \epsilon$ by (23). This together with (25) implies that $|\langle P_k h, g \rangle| \leq |k| \|h\| \|w\| < \epsilon |k| \|h\|$. Since ϵ is arbitrary, we have that $\|P_k h\| = 0$. Similarly, we can obtain that $\|P_0 h'\| = 0$ for every $h' \in R(A + kB)$. Therefore, $\|P_k - P_0\| = 0$. This completes the proof.

Remark 1. Theorem 1 is a generalization of Theorem 5.25 in [24] for operators to linear relations.

Next, we shall discuss the perturbations of a closed linear relation and its adjoint. We first recall a stability result of the closedness for linear relations.

Lemma 5 (Theorem 6.3 in [22]). *Let $T, S \in LR(X, Y)$ satisfy that $D(T) \subset D(S)$ and $S(0) \subset T(0)$. If S is T -bounded with T -bound less than 1, then $T + S$ is closed if and only if T is closed.*

Theorem 2. *Let $T, S \in LR(X, Y)$ satisfy that $D(T) \subset D(S)$ and $S(0) \subset T(0)$. Assume that T is closed and S is T -bounded with T -bound b . Further, if $b > 0$, set*

$$\Omega = \left\{ |z| < \frac{1}{b} : T + zS \text{ is closed} \right\}. \quad (29)$$

If $b = 0$, set

$$\Omega = \{ |z| \in \mathbb{C} : T + zS \text{ is closed} \}. \quad (30)$$

Then,

- (i) The set Ω is open.
- (ii) For every $z \in \Omega$, let Q_z denote the orthogonal projection from $X \times Y$ onto $T + zS$. Then, Q_z is continuous on Ω (with respect to the norm topology of $B(X \times Y)$, where $B(X \times Y)$ denote the bounded operators on $X \times Y$).

Proof

- (i) Suppose that T is closed and S is T -bounded with T -bound b . Set

$$\Phi := \{ r \geq 0 : \text{there exists } a \geq 0 \text{ such that } \|S(x)\| \leq a\|x\| + r\|T(x)\|, x \in D(T) \}. \quad (31)$$

Then, $b = \inf\{r : r \in \Phi\}$. Note that for every $z \in \mathbb{C}$, we have $D(T + zS) = D(T) \subset D(S)$ and $S(0) \subset (T + zS)(0) = T(0)$. Then, by Lemma 2, one has that

$$\|(T + zS)(x)\| \geq \|T(x)\| - \|(zS)(x)\| = \|T(x)\| - |z| \|S(x)\|. \quad (32)$$

This implies that

$$-|z| \|S(x)\| \leq \|(T + zS)(x)\| - \|T(x)\|. \quad (33)$$

Let $b > 0$. Given any $z_0 \in \Omega$. There is $\epsilon > 0$ such that $|z_0| < 1/(b + \epsilon)$. By the definition of infimum, there exists $r_1 \in \Phi$ such that $r_1 < b + \epsilon < 1/|z_0|$. Thus, $r_1 |z_0| < 1$. Further, there is a constant $a_1 \geq 0$ such that $\|S(x)\| \leq a_1 \|x\| + r_1 \|T(x)\|$, which together with (33) yields that

$$(1 - r_1 |z_0|) \|S(x)\| \leq a_1 \|x\| + r_1 \|(T + z_0 S)(x)\|. \quad (34)$$

If $b = 0$, for every $z_0 \in \Omega$, there exist $r_2 \in \Phi$ and $a_2 \geq 0$ such that $r_2 < 1/|z_0|$ and $\|S(x)\| \leq a_2 \|x\| + r_2 \|T(x)\|$. Again, by (33), we get that

$$(1 - r_2 |z_0|) \|S(x)\| \leq a_2 \|x\| + r_2 \|(T + z_0 S)(x)\|, \quad (35)$$

which together with (34) yields that S is $T + z_0 S$ -bounded. Hence, $T + zS$ is closed for z sufficiently near to z_0 by Lemma 5. Therefore, Ω is open.

- (ii) Define the linear relations $A, B \in LR(X, X \times Y)$ by

$$\begin{aligned} A(x) &= \{(x, f + z_0 g) : (x, f) \in T, (x, g) \in S\}, \\ B(x) &= \{(0, h) : (x, h) \in S\}, \end{aligned} \quad (36)$$

for every $x \in D(T)$. Obviously, $D(A) = D(B) = D(T)$, $A(0) = \{0\} \times T(0)$, and $B(0) = \{0\} \times S(0)$. Thus, $B(0) \subset A(0)$. Further, it is easy to verify that

$$\begin{aligned} \|A(x)\| &= \|(T + z_0 S)(x)\| + \|x\|, \\ \|B(x)\| &= \|S(x)\|, \end{aligned} \quad (37)$$

and

$$\begin{aligned} R(A) &= T + z_0 S, R(A + kB) \\ &= T + (z_0 + k)S. \end{aligned} \quad (38)$$

Note that S is $T + z_0 S$ -bounded. Then, there is a constant $c \geq 0$ such that $\|B(x)\| \leq c\|A(x)\|$. Therefore, Q_z is continuous on Ω by Theorem 1. The proof is complete.

In the following, we shall give a general perturbation result about a closed linear relation and its adjoint. \square

Lemma 6 (Theorem 4.30 in [4]). *Let M and N be closed subspace of X , and let P_M and P_N be the orthogonal projections onto M and N , respectively. Then, $M \perp N$ if and only if $P_M P_N = 0$ (or $P_N P_M = 0$) and if and only if $P_M + P_N$ is an orthogonal projection.*

Theorem 3. *Let $T, S \in LR(X, Y)$ satisfy that $D(T) \subset D(S)$ and $S(0) \subset T(0)$. Assume that T is closed, S is T -bounded*

with T -bound b , and S^* is T^* -bounded with T^* -bound b' . Let $r = \max\{b, b'\}$, if $r > 0$, set

$$\Omega = \left\{ |z| < \frac{1}{r} : T + zS, T^* + z^*S^* \text{ is closed} \right\}. \quad (39)$$

If $r = 0$, set

$$\Omega = \{ |z| \in \mathbb{C} : T + zS, T^* + z^*S^* \text{ is closed} \}. \quad (40)$$

Further, denote by Ω_0 the connected component of Ω that contains zero. Then,

$$(T + zS)^* = T^* + z^*S^*, \quad (41)$$

for all $z \in \Omega_0$.

Proof. Let Q_z and Q'_z be the orthogonal projections (in $X \times Y$) onto $T + zS$ and $U^{-1}(T^* + z^*S^*)$, respectively, where U is the flip-flop operator defined as (10). It follows from Theorem 2 that the operators Q_z and Q'_z depend continuously on $z \in \Omega$. By (11), we get that $T^\perp = U^{-1}T^*$. Hence, $X \times Y = T \oplus U^{-1}T^*$ and consequently,

$$Q_0 + Q'_0 = I. \quad (42)$$

By Proposition III.1.5 in [9], we have that $T^* + z^*S^* \subset (T + zS)^*$, which together with (11) implies that

$$U^{-1}(T^* + z^*S^*) \subset U^{-1}(T + zS)^* = T + zS, \quad z \in \Omega. \quad (43)$$

Thus, $Q_z Q'_z = Q'_z Q_z = 0$ for $z \in \Omega$ by Lemma 6. Again by Lemma 6, one has that $I - Q_z - Q'_z$ is an orthogonal projection for any $z \in \Omega$. Consequently, the value of $\|I - Q_z - Q'_z\|$ can be only 0 or 1. Since $\|I - Q_z - Q'_z\|$ depends continuously on $z \in \Omega$, which together with (42) yields that

$$\begin{aligned} \|I - Q_z - Q'_z\| &= \|I - Q_0 - Q'_0\| \\ &= 0, \quad z \in \Omega_0. \end{aligned} \quad (44)$$

Hence, $X \times Y = (T + zS) \oplus U^{-1}(T^* + z^*S^*)$. Therefore, $T^* + z^*S^* = U(T + zS) = (T + zS)^*$ for all $z \in \Omega_0$ by (11). This completes the proof.

If the relative bounds of S with respect to T and of S^* with respect to T^* are less than one, then $\{z \in \mathbb{C} : |z| \leq 1\} \subset \Omega_0$ by Lemma 5. Therefore, by Theorem 3, we have the following result. \square

Corollary 1. *Let $T, S \in LR(X, Y)$ satisfy that $D(T) \subset D(S)$ and $S(0) \subset T(0)$. Assume that T is closed, S is T -bounded with T -bound less than 1, and S^* is T^* -bounded with T^* -bound less than 1. Then, $(T + S)^* = T^* + S^*$.*

Remark 2. Let $T \in LR(X)$ be self-adjoint and $S \in LR(X)$ be Hermitian with $D(T) \subset D(S)$. Then, $S(0) \subset T(0)$ by Lemma 5.8 in [25]. If S is T -bounded with T -bound less than one, then S^* is T^* -bounded with T^* -bound less than one by Corollary III.1.13 in [17], and thus $(T + S)^* = T^* + S^* = T + S$ by Corollary 1, which means $T + S$ is self-adjoint. So, we shall remark that Corollary 1 generalizes

Theorem 5.2 in [25] for self-adjoint relations to general linear relations.

Remark 3. For studying stabilities of spectra of self-adjoint linear relations, we need to study stabilities of self-adjointness of linear relations. The results obtained in the present paper can be available in this case. Further, we shall apply these results to study stabilities of essential spectra of self-adjoint linear relations under some perturbations in our forthcoming paper.

Data Availability

No data were used to support this study.

Conflicts of Interest

The author declares that there are no conflicts of interest.

Acknowledgments

This research was supported by the National Natural Science Foundation of China (Grant no. 11790273).

References

- [1] S. Goldberg, *Unbounded Linear Operators, Theory and Applications*, McGraw-Hill, New York, NY, USA, 1966.
- [2] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, New York, NY, USA, 2nd edition, 1984.
- [3] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II: Fourier Analysis, Self-Adjointness*, Academic Press, Cambridge, CA, USA, 1972.
- [4] J. Weidmann, *Linear Operators in Hilbert Spaces Graduate Texts in Math* Vol. 68, Springer-Verlag, New York, NY, USA, 1980.
- [5] G. Ren and Y. Shi, "Self-adjoint extensions for discrete linear Hamiltonian systems," *Linear Algebra and Its Applications*, vol. 454, pp. 1–48, 2014.
- [6] Y. Shi and H. Sun, "Self-adjoint extensions for second-order symmetric linear difference equations," *Linear Algebra and Its Applications*, vol. 434, no. 4, pp. 903–930, 2011.
- [7] A. Favini and A. Yagi, "Multivalued linear operators and degenerate evolution equations," *Annali di Matematica Pura ed Applicata*, vol. 163, no. 1, pp. 353–384, 1993.
- [8] J. Behrndt, S. Hassi, and H. de Snoo, "Boundary value problems, wely functions, and differential operators," *Monographs in Mathematics*, vol. 108, 2020.
- [9] J. von Neumann, "Functional operator II: the geometry of orthogonal spaces," *Annals of Mathematics Studies*, vol. 22, 1950.
- [10] R. Arens, "Operational calculus of linear relations," *Pacific Journal of Mathematics*, vol. 11, no. 1, pp. 9–23, 1961.
- [11] T. Álvarez, A. Ammar, and A. Jeribi, "On the essential spectra of some matrix of linear relations," *Mathematical Methods in the Applied Sciences*, vol. 37, no. 5, pp. 620–644, 2014.
- [12] T. Y. Azizov, J. Behrndt, P. Jonas, and C. Trunk, "Compact and finite rank perturbations of closed linear operators and relations in Hilbert spaces," *Integral Equations and Operator Theory*, vol. 63, no. 2, pp. 151–163, 2009.
- [13] A. G. Baskakov and A. S. Zagorskii, "Spectral theory of linear relations on real Banach spaces," *Mathematical Notes*, vol. 81, no. 1-2, pp. 15–27, 2007.

- [14] E. Chafai and M. Mnif, "Perturbation of normally solvable linear relations in paracomplete spaces," *Linear Algebra and Its Applications*, vol. 439, no. 7, pp. 1875–1885, 2013.
- [15] E. A. Coddington, "Extension theory of formally normal and symmetric subspaces," *Memoirs of the American Mathematical Society*, vol. 134, 1973.
- [16] E. A. Coddington and A. Dijksma, "Self-adjoint subspaces and Eigen function expansions for ordinary differential subspaces," *Journal of Differential Equations*, vol. 20, no. 2, pp. 473–526, 1976.
- [17] R. Cross, "Multivalued linear operators," *Monographs and Textbooks in Pure and Applied Mathematics*, vol. 213, 1998.
- [18] R. Cross, A. Favini, and Y. Yakubov, "Perturbation results for multivalued linear operators," *Parabolic Problems*, vol. 80, pp. 111–130, 2011.
- [19] A. Dijksma and H. de Snoo, "Self-adjoint extensions of symmetric subspaces," *Pacific Journal of Mathematics*, vol. 54, no. 1, pp. 71–100, 1974.
- [20] Y. Shi, C. Shao, and Y. Liu, "Resolvent convergence and spectral approximations of sequences of self-adjoint subspaces," *Journal of Mathematical Analysis and Applications*, vol. 409, no. 2, pp. 1005–1020, 2014.
- [21] Y. Shi, C. Shao, and G. Ren, "Spectral properties of self-adjoint subspaces," *Linear Algebra and Its Applications*, vol. 438, no. 1, pp. 191–218, 2013.
- [22] Y. Shi, G. Xu, and G. Ren, "Boundedness and closedness of linear relations," *Linear and Multilinear Algebra*, vol. 66, no. 2, pp. 309–333, 2018.
- [23] H. Sun, Q. Kong, and Y. Shi, "Essential spectrum of singular discrete linear Hamiltonian systems," *Mathematische Nachrichten*, vol. 289, no. 2-3, pp. 343–359, 2016.
- [24] H. Sun and Y. Shi, "Spectral properties of singular discrete linear Hamiltonian systems," *Journal of Difference Equations and Applications*, vol. 20, no. 3, pp. 379–405, 2014.
- [25] G. Xu and Y. Shi, "Perturbations of spectra of closed subspaces in Banach spaces," *Linear Algebra and Its Applications*, vol. 531, pp. 547–574, 2017.
- [26] G. Xu and Y. Shi, "Essential spectra of self-adjoint relations under relatively compact perturbations," *Linear and Multilinear Algebra*, vol. 66, no. 12, pp. 2438–2467, 2018.